Big-O notation

Lecture 10

Running time of selection sort

- We showed that running selection sort on an array of n elements takes in the worst case $T(n) = 1 + 15\,n + 5\,n^2$ primitive operations
- When n is large, $T(n) \approx 5\,n^2$
- When n is large,

  $T(2n) / T(n) \approx 5\,(2n)^2 / 5\,n^2$

  ≈ 4

  Doubling n quadruples T(n)

  N.B. That is true for any coefficient of n^2 (not just 5)

n	T(n)
10	661
20	2301
30	4951
40	8601
...	...
1000	5015001
2000	20030001

Big - O notation

- Goals:
  - Simplify the discussion of algorithm running times
  - Describe how the running time of an algorithm increases as a function of n (the size of the problem), when n is LARGE
  - Get rid of terms that become insignificant when n is large
- We will say things like:

  The worst-case running time of selectionSort on an array of n elements is $O(n^2)$

  The worst-case running time of mergeSort on an array of n elements is $O(n \log(n))$

Big-O definition

- Let f(n) and g(n) be two non-negative functions defined on the natural numbers N
- We say that f(n) is O(g(n)) if and only if:
  - There exists an integer n_0 and a real number c such that: for all $n \geq n_0$, $f(n) \leq c \bullet g(n)$

  More mathematically, we would write
  - $\exists\, n_0 \in N, \exists\, c \in R : \forall\, n \geq n_0, f(n) \leq c \bullet g(n)$
- N.B. The constant c must *not* depend on n

Intuition and visualization

- "f(n) is O(g(n))" iff there exists a point n_0 beyond which f(n) is less than some fixed constant times g(n)



$f(n) = 5 + 3\,n^2$

$g(n) = 1 + n^2$



f(n) is O(g(n)), because there exists $n_0 = 2$ and c = 4 such that for all $n \geq n_0$, $f(n) \leq c \bullet g(n)$

Proving big-O relations

- To prove that f(n) is O(g(n)), we must find n_0 and c such that $f(n) \le c \bullet g(n)$
- Example: Prove that $5 + 3n^2$ is $O(1 + n^2)$

We need to pick c greater 3. Let's pick c = 5.

If we choose $n_0 = 1$, we get that if $n \ge n_0$, then

$$5 + 3n^2 \le 5 + 5n^2 \quad \text{(since } n \ge n_0\text{)}$$
$$= 5(1 + n^2)$$
$$= c(1 + n^2)$$

Examples

- Prove that 2n + 3 is O(n)

Examples

- Prove that $f(n) = 10^{100}$ is O(1)

Examples

- Prove that n (sin(n) + 1) is O(n)

Proving that f(n) is *not* O(g(n))

- To prove that f(n) is *not* O(g(n)), one must show that for any n_0 and c, there exists an $n \ge n_0$ such that $f(n) > c\, g(n)$
- Procedure: Assume n_0 and c are given, and find a value of n such that $f(n) > c\, g(n)$. The value of n will usually depend on n_0 and c

Examples

- Prove that n^2 is *not* O(n)

Examples

- Prove that n (sin(n) + 1) is O(n)

Examples

- Prove that 2^n is *not* $O(n^3)$