

(1)

From DFA to regular expressions over Σ :

Thm

For a DFA $M = (S, s_0, \delta, F)$ there is a regular expression r s.t. $L_M = \underline{\text{Lr}}$

Proof

A DFA has a finite number of states, say n . Number the states 1 through n . We are going to define a family of regular expressions $R_{ij}^{(k)}$ where $i, j, k \in \{1, 2, \dots, n\}$. The meaning of $R_{ij}^{(k)}$ is the regular expression describing all words w such that if M is in state i , when M reads w it will end up in state j and all the states along the way are numbered k or less. We will construct this starting from $k=0$ & go up to $k=n$.

$k=0$ There are no states numbered 0 or less.

This means there should be a direct path from i to j if $i=j$ there is a length 0 path from i to itself.

If $\exists a \in \Sigma$ s.t. $\delta(s_i, a) = s_j$ we set

$$R_{ij}^{(0)} = a$$

If there are several such letters in Σ , say a_1, \dots, a_ℓ

$$R_{ij}^{(0)} = a_1 + a_2 + \dots + a_\ell; \text{ where } \forall a_m, \delta(s_i, a_m) = s_j.$$

If $i=j$ we do exactly the same except we add ϵ

$$R_{ii}^{(0)} = \epsilon + a_1 + \dots + a_\ell \text{ where for each } a_m, \delta(s_i, a_m) = s_i.$$


Suppose we have constructed all the regular expressions for every i, j, k for k up to some value.

Now consider $R_{ij}^{(k+1)}$:

$$R_{ij}^{(k+1)} = R_{ij}^{(k)} + R_{ik}^{(k)} \left(R_{kk}^{(k)} \right)^* R_{kj}^{(k)}$$

How did we get this?

(2)




This picture makes clear why. The $R_{ij}^{(k)}$ term represents the paths we already had. Now we need to add new paths that use the node $(k+1)$. To get from i to $(k+1)$ we can travel along any path in $R_{i(k+1)}^{(k)}$. We can return to $(k+1)$ several (or zero) times, hence the $(R_{(k+1)(k+1)}^{(k)})^*$ and then we must get to j using $R_{(k+1)j}^{(k)}$

Clearly all the constructs we are using give us regular expressions. When we get $R_{ij}^{(n)}$ for all i, j we can now construct the regular expression for L_M as follows. Let the start state have number 1 (we are free to choose the numbering) and let the final states have numbers i_1, \dots, i_p . Then

$$L_M = R_{1i_1}^{(n)} + R_{1i_2}^{(n)} + \dots + R_{1i_p}^{(n)} \blacksquare$$

Example



$$\begin{aligned}
 R_{11}^{(0)} &= \epsilon + b & R_{12}^{(0)} &= a & R_{21}^{(0)} &= \phi & R_{22}^{(0)} &= \epsilon + a + b \\
 R_{11}^{(1)} &= R_{11}^{(0)} + R_{11}^{(0)} (R_{11}^{(0)})^* R_{11}^{(0)} = b^*; & R_{12}^{(1)} &= b^* a & R_{21}^{(1)} &= \phi & R_{22}^{(1)} &= \epsilon + a + b \\
 R_{11}^{(2)} &= b^*; & R_{12}^{(2)} &= b^* a + b^* a (\epsilon + a + b)^* (\epsilon + a + b) = b^* a (a + b)^* & R_{22}^{(2)} &= (a + b)^* \\
 R_{12}^{(2)} & \text{ is } L_M
 \end{aligned}$$